

## CURRICULUM VITAE



### PERSONAL INFORMATION

NAME  
ADDRESS  
TELEPHONE  
PEC  
E-mail  
  
NATIONALITY  
  
DATE OF BIRTH

**Albertin Andrea**

[albertinsimone@pec.it](mailto:albertinsimone@pec.it)

[albertinsimone@yahoo.it](mailto:albertinsimone@yahoo.it)

**Italian**

**MAY 22 1965**

Veteran professional with more than 20 years of experience as an anaesthetist. An effective educator with about 10 years experience in corporate training. Major interests include clinical pharmacokinetics and pharmacodynamics, drug administration technology, target controlled infusion techniques, regional anaesthesia and airway management.

### WORK EXPERIENCE

**1<sup>st</sup> Apr 2021 – Freelance**

**1<sup>st</sup> Feb 2013 – 1<sup>st</sup> Mar 2021**

**1<sup>st</sup> Feb 2010 - 25<sup>th</sup> Mar 2021**

**1<sup>st</sup> Jan 2008 - 31<sup>st</sup> Jan 2010**

**3<sup>rd</sup> May 1999 – 31<sup>st</sup> Dec 2007**

**1<sup>st</sup> Oct 1998 – 28<sup>th</sup> Feb 1999**

**1<sup>st</sup> Jan 1995 - 30<sup>th</sup> Sept 1998**

**1<sup>st</sup> Aug 1994 – 31<sup>st</sup> Dec 1994**

*Head Anaesthetist St Mary Clinic - Castellanza (VA) Italy*

*Head Anaesthetist San Giuseppe Hospital – Milan Italy*

*Deputy Chief Anaesthetist IRCCS Multimedica Sesto San Giovanni*

*Anaesthetist San Raffaele Hospital Milan*

*Anaesthetist Maggiore Hospital Novara as trainee*

*Anaesthetist Magenta Hospital*

*Anaesthetist Voghera Hospital*

### EDUCATION AND TRAINING

**From January 2014 up to January 2022**

**2012-Today**

**2005-2007**

National certification for Associate Professor

External Professor in Anaesthesia and Rianimation – University of Milan. I School of Specialization (Prof. N. Stocchetti)

Head of Clinical Research Area San Raffaele Hospital Milan

- 2001-2007** External Professor in Anaesthesia and Rianimation – Life and Health University San Raffaele Milan (Prof. G. Torri)
- 1999-2001** External Professor in Anaesthesia and Rianimation – University of Milan. II School of Specialization (Prof. G. Torri)
- 1993** Academic Degree in Medicine and Surgery Pavia University 110/110  
Specialization Degree in Anaesthesia and Rianimation 70/70

## COMPETENCIES TO RELATIONSHIPS

Futus corporation future systems COURSE: "RHETORIC"

Futus corporation future systems COURSE: "THE JOY TO SELL"

Futus corporation future systems COURSE: "THE CHALLENGE OF LEADER"

## COMPETENCIES IN ORGANIZATION

Head Clinical Research Area  
San Raffaele Hospital Milan  
Research Areas

**2005-2007**

Mechanical ventilation and anaesthesia  
Inhalational anaesthetics  
Regional Anaesthesia  
Total intravenous anaesthesia  
Techniques of sedations and analgesia  
Bariatric anaesthesia  
Devices for intravenous anaesthetics infusion  
Devices for the monitoring of anaesthesia depth

**2010 – 2012**

*Head physician San Giuseppe  
Multimedica Hospital – Milan*

Management training course for heads of hospital department – Lombardia

## OCCUPATIONAL SKILLS AND COMPETENCES

**Managerial Skills**

Resource Management; Staff Management; Personnel Management

<i>Functional Skills</i>	<p>General Anaesthesia to cover a wide range of surgical specialities</p> <p>Total Intra Venous Anaesthesia</p> <p>Spinal and Epidural Procedures</p> <p>Loco regional anaesthesia</p> <p>Post Anaesthesia Care</p> <p>Medical Training and Education</p> <p>Drug Management</p> <p>Team Management</p>
<b>FIRST LANGUAGE</b>	<b>Italian</b>
<b>OTHER LANGUAGES</b>	<p><b>ENGLISH</b></p> <p><b>FRANCAIS</b></p>
<b>MEDICAL AFFILIATIONS</b>	<p>SIAARTI member</p> <p>AAROI member</p> <p>Foundation Member of World Society of Intravenous Anaesthesia</p>

## CLINICAL TRIALS

### Indexed Scientific Articles = 44

1. Cappelleri G., Ghisi D., Fanelli., **Albertin A.**, Somalvico F., Adegheri G. Does continuous sciatic nerve block improve postoperative analgesia and early rehabilitation after total knee arthroplasty? A prospective randomized, double blind study. *Reg Anesth Pain Med* 2011 Sept.-Oct 36(5); 489-492.
2. La Colla L, **Albertin, A**, La Colla, G, Porta A, Aldegheri G, Mangano A, Di Candia D, Gigli F. No adjustment vs "fictitious height" as height input for remifentanil Target Controlled Infusion (TCI) in morbidly obese patients: a preliminary, pilot study on 15 female patients. *Clin Pharmacokinet*, 2010;49(2):131-9.
3. Cappelleri G., Ghisi D., Fanelli A., Aldegheri G., La Colla L., **Albertin A.** Posterior psoas vs 3-in-1 approach for lateral femoral cutaneous and obturator nerve block for anterior cruciate ligament repair. *Minerva Anesthesiol.* 2009 Oct;75(10):568-73.
4. La Colla L., **Albertin A.**, La Colla G. Pharmacokinetic model-driven remifentanil administration in the morbidly obese: the "critical weight" and the "fictitious height". A possible solution to an unsolved problem? *Clin Pharmacokinet.* 2009;48(6):397-8.
5. La Colla L, **Albertin A**, La Colla G, Ceriani V, Lodi T, Porta A, Aldegheri G, Mangano, A, Khairallah, I, Fermo, I. No adjustment vs. adjustment formula as input weight for propofol Target Controlled Infusion (TCI) in morbidly obese patients. *Eur J Anaesthesiol.*, 2009;26(5):362-9
6. **Albertin A**, La Colla L, Gandolfi A, Colnaghi E, Mandelli D, Gioia G, Fraschini G. Greater peripheral blood flow but less bleeding with propofol vs sevoflurane during spine surgery: a possible physiologic model? *Spine* 2008;33(18):2017-22.
7. Dedola E, **Albertin A**, Poli D, La Colla L, Gandolfi A, Martani C, Torri G. Effect of nitrous oxide on desflurane MACBAR at two target-controlled concentrations of remifentanil. *Minerva Anesthesiol* 2008 May;74(5):165-72
8. La Colla L, Poli D, **Albertin A**, La Colla G, Mangano A. Preoperative gum chewing may increase gastric fluid retention. Time to change our minds and review current guidelines? *Obes Surg* 2008 Apr;18(4):430-1.
9. La Colla L., La Colla G., **Albertin A.**, Poli D., Baruffali Preis FW, Mangano A. The use of propofol and remifentanil for the anaesthetic management of a super obese patient. *Anaesthesia* 2007 Aug; 62(8): 842-5.
10. La Colla L., **Albertin A.**, La Colla G., Mangano A. Faster wash in and recovery for desflurane vs sevoflurane in morbidly obese patients when no premedication is used. *Br J Anaesth* 2007 Sep; 99(3): 353-8
11. La Colla G., La Colla L., Turi S., Poli D., **Albertin A.**, Pasculli N., Bergonzi PC., Gonfalini M., Ruggirei F. Effect of morbid obesity on kinetic of desflurane: wash-in wash-out curves and recovery times. *Minerva Anesthesiol* 2007 May; 73(5): 275-9.
12. La Colla L., La Colla G., Poli D., **Albertin A.** A new twist in approaching difficult tracheal intubation. Description of a economic, easy, and effective way: the sigma way. *Anesthesiology* 2007 May; 106(5): 1069-70
13. Mattioli C., Gemma M., Baldoli C., Sessa M., **Albertin A.**, Beretta L. Sedation for children with metachromatica leukodystrophy undergoing MRI *Paediatric Anaesth* 2007 Jan; 17(1): 64-9
14. **Albertin A.**, Poli D., La Colla L., Gonfalini M., Turi S., Pasculli N., La Colla G.,

Bergonzi P., Dedola E., Fermo I.  
Predictive performance of "Servin's formula" during BIS-guided propofol-remifentanil target-controlled infusion in morbidly obese patients  
Br J Anaesth 2007 Jan; 98(1): 66-75

15. **Albertin A.**, La Colla G., La Colla L., Bergonzi P., Deni F., Moizo E.  
Effect site concentrations of remifentanil maintaining cardiovascular homeostasis in response to surgical stimuli during bispectral index guided propofol anesthesia in seriously obese patients.  
Minerva Anesthesiol 2006 Nov; 72(11): 915-24.

16. Fanti L., Agostoni M., Arcidiano PG., **Albertin A.**, Strini G., Carrara S., Guslandi M., Torri G., Testoni PA.  
Target-controlled infusion during monitored anesthesia care in patients undergoing EUS: propofol alone versus midazolam plus propofol. A prospective double-blind randomized controlled trial.  
Dig Liver Dis 2007 Jan; 39(1): 81-6

17. **Albertin A.**, Dedola E., Bergonzi P., Lombardo F., Fusco T., Torri G.  
The effect of adding two target-controlled concentrations (1-3 ng/ml) of remifentanil on MAC BAR of desflurane.  
Eur J Anaesthesiol 2006 Jun; 23(6): 510-6.

18. **Albertin A.**, Marchetti C., Mamo D., Poli D., Dedola E.  
Intracranial hypotension: a case of spontaneous arachnoid rupture in a parturient.  
Anesthesiology 2006 Mar; 104(3): 609-10.

19. **Albertin A.**, Casati A., Bergonzi P., Moizo E., Lombardo F., Torri G.  
The effect of adding nitrous oxide on MAC of sevoflurane combined with two target-controlled concentrations of remifentanil in women.  
Eur J Anaesthesiol 2005 Jun; 22(6): 431-7

20. **Albertin A.**, Casati A., Lombardo F., Valeri R., Travaglino V., Bergonzi P., Torri G.  
The effect-site concentration of remifentanil blunting cardiovascular responses to tracheal intubation and skin incision during bispectral index-guided propofol anesthesia.  
Anesth Analg 2005 Jul; 101(1): 125-30.

21. Moizo E., Berti M., Marchetti C., Deni F., **Albertin A.**, Muzzolon F., Antonino A.  
Acute Pain Service and multimodal therapy for postsurgical pain control: evaluation of protocol efficacy.  
Minerva Anesthesiol 2004 Nov; 70(11): 779-87

22. **Albertin A.**, Casati A., Bergonzi P., Fano G., Torri G.  
Effect of two target-controlled concentrations (1 and 3 ng/ml) of remifentanil on MAC(BAR) of sevoflurane.  
Anesthesiology 2004 Feb; 100(2): 255-9

23. Danelli G., Berti M., Casati A., **Albertin A.**, Deni F., Nobili F., Torri G.  
Spinal block or total intravenous anaesthesia with propofol and remifentanil for gynaecological outpatient procedures.  
Eur J Anaesthesiol 2002 Aug; 19(8): 594-9

24. Danelli G., Berti M., Perotti V., **Albertin A.**, Baccari P., Deni F., Fanelli G., Casati A.  
Temperature control and recovery of bowel function after laparoscopic or laparotomic colorectal surgery in patient receiving combined epidural/general anesthesia and postoperative epidural analgesia  
Anesth Analg 2002 Aug; 95(2): 467-71

25. Berti M., Danelli G., **Albertin A.**, Deni F., Moizo E., Muzzolon F.  
Systemic analgesia after peripheral blocks  
Minerva Anesthesiol 2001 Sep;67(9) suppl 1:195-9. Review
26. Torri G., Casati A., **Albertin A.**, Comotti L., Bignami E., Scarioni M., Paganelli M.  
Randomized comparison of isoflurane and sevoflurane for laparoscopic gastric banding in morbidly obese patients.  
J Clin Anesth 2001 Dec;13(8):565-70
27. Casati A., **Albertin A.**, Danelli G., Deni F., Scarioni M., Santorsola R., Nucera M.  
Implementino sevoflurane anesthesia with small doses opioid for upper abdominal surgery. Postoperative respiratory function after either remifenatnil or fentanyl.  
Minerva Anesthesiol 2001 Sep;67(9):621-8
28. Di Benedetto P., Bertini L., Casati A., Borghi B., **Albertin A.**, Fanelli G.  
A new posterior approach to the sciatic nerve block: a prospective, randomized comparison with the classic posterior approach.  
Anesth Analg 2001 Oct;93(4):1040-4
29. Casati A., Fanelli G., Beccaria P., Magistris L., Albertin A., Torri G. The effects of single or multiple injections on the volume of 0.5% ropivacaine required for femoral nerve blockade.  
Anesth Analg 2001 Jul;(93(1):183-6.
30. Fanelli g., Casati A., Magistris L., Berti M., **Albertin A.**, Scarioni M., Torri G.  
Fentanyl does not improves the nerve block characteristics of axillary brachial plexus anaesthesia performed with ropivacaine.  
Acta Anaesthesiol Scand 2001 May;45(5):590-4
31. Casati A., Fanelli G., **Albertin A.**, Deni F., Danelli G., Grifoni F., Torri G.  
Small doses of remifentanil or sufentanil for blunting cardiovascular changes induced by tracheal intubation: a double blind comparison.  
Eur J Anaesthesiol 2001 Feb;18(2):108-12
32. Fanelli G., Casati A., Beccarla P., Cappelleri G., **Albertin A.**, Torri G.  
Interscalene brachial plexus anaesthesia with small volumes of ropivacaine 0.75%: effect of the injection technique on the onset time of nerve blockade.  
Eur J Anaesthesiol 2001 Jan;18(1):54-8.
33. **Albertin A.**, Casati A., Deni F., Danelli G., Comotti L., Grifoni F., Fanelli G.  
Clinical comparison of either small doses of fentanyl or remifentanil for blunting cardiovascular changes induced by tracheal intubation.  
Minerva Anesthesiol. 2000 Oct;66(10):69-6
34. Casati A., **Albertin A.**, Fanelli G., Deni F., Berti M., Danelli G., Grifoni F., Torri G.  
A comparison of remifentanil and sufentanil as adjuvants during sevoflurane anesthesia with epidural analgesia for upper abdominal surgery: effects on postoperative recovery and respiratory function.  
Anesth Analg 2000 Nov;91(5):1269-73
35. Berti M., **Albertin A.**, Casati A., Palmisano S., Municino G., Da Gama Malcher M., De Ponti A. A prospective, randomized comparison of dexketoprofen, ketoprofen or paracetamol for postoperative analgesia after outpatient knee arthroscopy. Minerva Anesthesiol 2000 Jul-Aug;66(7-8):549-54
36. Berti M., Danelli G., **Albertin A.**, Casati A., Cucchi C., Torri G.  
Bispectral Index: clinical effectiveness and role in reducing anesthetic drug consumption  
Minerva Anesthesiol. 2000 May;66(5):394-7. Review

37. Berti M., Casati A., Fanelli G., **Albertin A.**, Palmisano S., Danelli G., Comotti L., Torri G.  
0.2% ropivacaine with or without fentanyl for patient-controlled epidural analgesia after major abdominal surgery: a double blind study.  
J Clin Anesth 2000 Jun;12(4):292-7
38. Magistris L., Casati A., **Albertin A.**, Deni F., Danelli G., Borghi B., Fanelli G.  
Combined sciatic-femoral nerve block with 0.75% ropivacaine: effects of adding a systemically inactive dose of fentanyl.  
Eur J Anaesthesiol 2000 Jun;17(6):348-53.
39. Cappelleri G., Casati A., Fanelli G., Borghi B., Anelati D., Berti M., **Albertin A.**  
Unilateral spinal anesthesia or combined sciatic-femoral nerve block for day case arthroscopy. A prospective, randomized, comparison.  
Minerva Anesthesiol. 2000 Mar;66(3):131-6; discussion 137
40. Casati A., Fanelli G., **Albertin A.**, Deni F., Anelati D., Antonino A., Beccaria P.  
Interscalene brachial plexus anesthesia with either 0.5% ropivacaine or 0.5% bupivacaine.  
Minerva Anesthesiol. 2000 Jan-Feb;66(1-2):39-44.
41. Berti M., Fanelli., Casati A., **Albertin A.**, Palmisano S., Deni F., Perotti V., Torri G.  
Patient supplemented epidural analgesia after major abdominal surgery with bupivacaine/fentanyl or ropivacaine/fentanyl.  
Can J Anaesth 2000 Jan;47(1):27-32.
42. Casati A., Cappelleri G., Fanelli G., Magistris L., Beccaria P., **Albertin A.**, Torri G.  
The pressor response after laryngeal mask or cuffed oropharyngeal airway insertion.  
Acta Anaesthesiol Scand 1999 Nov;43(10):1053-6
43. Casati A., Fanelli G., Cappelleri G., **Albertin A.**, Anelati D., Magistris L., Torri G.  
Arterial to end-tidal carbon dioxide tension difference in anaesthetized adults mechanically ventilated via laryngeal mask or a cuffed oropharyngeal airway.  
Eur J Anaesthesiol. 1999 Aug;16(8):534-8.
44. Berti M., **Albertin A.**, Palmisano S. Consequences of perioperative hypothermia  
Minerva Anesthesiol. 1999 May;65(5suppl):42-5. Review

## Letters to the Editor N=22

1. La Colla L, Mangano A, Mangano A, **Albertin A**. Effects of nonpathogenic gram-negative bacterium *Vitreoscilla filiformis* lysate on atopic dermatitis: a prospective, randomized, double-blind, placebo-controlled clinical study: does this make a real difference? *Br J Dermatol.*, 2009 Aug; 161(2): 477-8.
2. Mangano A, **Albertin A**, La Colla L. Clinical investigation of the novel iron-chelating agent, CP94, to enhance topical photodynamic therapy of nodular basal cell carcinoma: when statistics makes difference. *Br J Dermatol.*, 2009 Jul; 161(1): 215-6.
3. La Colla L, Mangano A, **Albertin A**. Continuous infusion of intraperitoneal bupivacaine after laparoscopic surgery: a randomized controlled trial. What about statistical power and analysis? *Obes Surg*, 2009 Jun;19(6):817-8
4. Mangano A, **Albertin A**, La Colla L. Patient controlled sedation using a standard protocol for dressing changes in burns: patients' preference, procedural details and a preliminary safety evaluation. Are studies always adequately powered and analyzed? *Burns*, 2010 Sep; 36(6): 948.
5. La Colla L, Mangano A, **Albertin A**. Beneficial effects of humidified, warmed carbon dioxide insufflation during laparoscopic bariatric surgery: a randomized clinical trial. What if sample size calculation made difference? *Obes Surg*, 2010 Dec; 20(12): 1747.
6. La Colla L, **Albertin A**, Mangano A, Mangano A. Effects of preoperative local ropivacaine infiltration on postoperative pain scores in infants and small children undergoing elective cleft palate repair: statistical considerations to transform interesting hypotheses into significant conclusions. *J Craniof Surg* 2009 Sep; 20(5): 1627-8
7. Mangano A, **Albertin A**, La Colla L. A double-blind, randomized, placebo-controlled, two-dose comparative study of botulinum toxin type A for treating glabellar lines in Japanese subjects: what if sample size and statistical tests mattered? *Aesthetic Plast Surg*, 2009 Sep; 33(5): 788.
8. Mangano A, **Albertin A**, Mangano A, Mangano C, La Colla L. Comparison between the use of an ultrasonic tip and a microhead handpiece in periradicular surgery: A prospective randomised trial. What about study design and statistics? *Br J Oral Maxillofac Surg*, 2009 Jun;47(4):332
9. La Colla L, **Albertin A**, Mangano A, Mangano A. Progressive tension sutures in the prevention of postabdominoplasty seroma: a prospective, randomized, double-blind clinical trial: the quest for statistical significance goes through sample size calculation. *Plast Reconstr Surg*, 2009 Jul; 124(1): 333-1.
10. Mangano A, **Albertin A**, Mangano A, La Colla L. The use of suction drains in abdominal dermolipectomy: a randomized clinical trial: are "significant" findings always so? *Plast Reconstr Surg*, in press.
11. Mangano A, **Albertin A**, Mangano A, La Colla L. Temporal hollowing following coronal incision: a prospective, randomized, controlled trial: statistics is crucial to draw adequate conclusions in plastic surgery. *Plast Reconstr Surg*, 2009 Aug; 124(2) 669-70



12. La Colla L, Mangano A, Mangano A, **Albertin A**. Dipyron increases the blood flow of arterial dorsal skin flaps: are prospective studies always properly designed? *Aesthetic Plast Surg*, 2009 Mar;33(2):257.
13. Mangano A, **Albertin A**, La Colla L. Use of 2-octyl-cyanoacrylate skin adhesive (Dermabond) for wound closure following reduction mammoplasty: a prospective, randomized intervention study: "Tips and Tricks" to improve statistical analysis and significance of results. *Plast Reconstr Surg*, 2009 Aug; 124(2): 669.
14. La Colla L, Mangano A, **Albertin A**. A Prospective, Randomized, Double-Blind, Controlled Trial of Continuous Local Anesthetic Infusion in Cosmetic Breast Augmentation: what lies beyond data? *Plast Reconstr Surg*, 2009 May; 123(5): 1636-7.
15. La Colla L, Mangano A, Mangano A, **Albertin A**. Prospective, Randomized, Double-Blind Trial of Local Anesthetic Infusion and Intravenous Narcotic Patient-Controlled Anesthesia Pump for Pain Management after Free TRAM Flap Breast Reconstruction: are we on the right track? *Plast Reconstr Surg*, 2009; May; 123(5): 362-9.
16. Mangano A, **Albertin A**, La Colla L. Povidone Iodine versus Chlorhexidine in Skin Antisepsis before Elective Plastic Surgery Procedures: A Randomized Controlled Trial. Is statistical correctness always pursued? *Plast Reconstr Surg*, 2009 Jul; 124(1): 340.
17. Mangano A, **Albertin A**, La Colla L. Patient controlled sedation using a standard protocol for dressing changes in burns: patients' preference, procedural details and a preliminary safety evaluation. Are studies always adequately powered and analyzed? *Burns*, 2010 Sep; 36(6): 948.
18. La Colla L, Mangano A, Mangano A, **Albertin A**. Effects of nonpathogenic gram-negative bacterium *Vitreoscilla filiformis* lysate on atopic dermatitis: a prospective, randomized, double-blind, placebo-controlled clinical study: does this make a real difference? *Br J Dermatol*, 2009 Aug;161(2): 477-8.
19. La Colla L, Poli D, **Albertin A**, La Colla G, Mangano A. Preoperative gum chewing may increase gastric fluid retention. Time to change our minds and review current guidelines? *Obes Surg*. 2008;18:430-31.
20. La Colla L, La Colla G, Poli D, **Albertin A**. A new twist in approaching difficult tracheal intubation. Description of an economic, easy and effective way: the Σ way. *Anesthesiology* 2007; 106:1069-70.
21. **Albertin A**, Marchetti C., Mamo D., Poli D., Dedola E. Contrast imaging while breast-feeding *Anesthesiology* 2006 Dec; 105(6): 1277 replay
22. La Colla L., **Albertin A**. No difference in blood loss during endoscopic sinus surgery with total intravenous anesthesia? *JAMA Otolaryngol Head Surg* 2013 Oct; 139(10): 1077.

**Non indexed Abstracts = 3**

1. Early labour epidural analgesia in nulliparous  
IV Italian National Meeting ESRA S. Margherita Ligure 19-22 Giugno 1996
2. Incremento dell'utilizzo del blocco neuroassiale in anestesia ostetrica  
International Joint Meeting of Obstetric e Pediatric Anaesthesia (CIAO e SARNEPI)  
Abano T. October 25-26 1996
3. Conseguenze ostetriche dell'analgesia epidurale precoce nel travaglio iniziale  
International Joint Meeting of Obstetric e Pediatric Anaesthesia (CIAO e SARNEPI)  
Abano T. October 25-26 1996

## Indexed Abstracts = 71

1. Total Intravenous Anaesthesia (TIVA) with Remifentanil and Propofol for Elective Intraabdominal Surgery  
SIAARTI Udine 1998. Minerva Anesthesiol vol. 2 1998
2. PCA per il controllo del dolore postoperatorio dopo TIVA con Remifentanil in chirurgia addominale  
VIII Congresso Nazionale Società Italiana dei Clinici del Dolore. Villa Erba-Cernobbio 26-28 Novembre 1998
3. Blocco del plesso celiaco nell'interventistica delle vie biliari  
ESRA 6<sup>th</sup> Italian Charter Meeting Milano 6-8 Dicembre 1998
4. Sevoflurane pharmacokinetic in morbidly obese patients  
SIAARTI Napoli 2000. Minerva Anesthesiol vol.66 2000
5. Sevoflurane balanced general anesthesia with small doses fentanyl or remifentanil: early effects on recovery and respiratory function  
SIAARTI Napoli 2000. Minerva Anesthesiol vol.66 2000
6. PCEA with 0.2% ropivacaine with or without small doses sufentanil after arthroscopic cruciate ligament repair  
SIAARTI Napoli 2000. Minerva Anesthesiol vol.66 2000
7. BIS monitoring in operative hysteroscopy  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
8. Improving postoperative analgesia after axillary brachial plexus anaesthesia with 0.75% ropivacaine: a double blind evaluation of adding clonidine  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
9. 0.75% ropivacaine and fentanyl for axillary brachial plexus block  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
10. Effects of multiple injections on volume of 0.5% ropivacaine required for femoral nerve blockade  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
11. General anaesthesia allow a fast sciatic discharge than combined sciatic femoral nerve  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
12. Spinal block or total intravenous anaesthesia for outpatient hysteroscopy  
ESA 2001 Gothenburg 2001. Eur J Anaesthesiol. Vol 18 Suppl 21
13. Fentanyl 2  $\mu$ g mL<sup>-1</sup> reduces ropivacaine consumption during patient-controlled epidural analgesia after major abdominal surgery. Eur J Anaesth 2000;19:A594.
14. Single or multiple injections for interscalene brachial plexus anesthesia with 0.75% ropivacaine? Anesthesiology 2000;93:A814.
15. Ropivacaine and bupivacaine equipotency in peripheral nerve blocks. Minerva Anesthesiol 2000
16. Spinal block or total intravenous anesthesia with propofol and remifentanil for gynecological outpatient procedures. Anesthesiology 2001;95:A460.
17. Laparoscopic and laparotomic, colorectal surgery in patients receiving general-epidural anesthesia. Eur J Anaesthesiol 2002;19:A-14.
18. The effect of remifentanil on sevoflurane requirements for somatic and sympathetic responses to surgical incision. Minerva Anesthesiol 2002;68(S.2):247.

19. Effect of nitrous oxide on combined sevoflurane Remifentanil MACbar in women  
SIAARTI Roma 2003 Minerva Anesthesiol vol 69 suppl 2 N 9 Sept 2003
20. Effect of thoracic epidural block on oxygenation during one-lung ventilation: a pilot study. Reg Anesth Pain Med 2003;28:A28.
21. Farmacoeconomia in anesthesia: alcune considerazioni di ordine generale  
XIII Convegno Sicurezza in Anestesia – Napoli, 12-13 Dicembre 2003
22. Patient – controlled epidural analgesia after anterior cruciate ligament repair: is it useful adding sufentanil to 0.2% ropivacaine?  
Eur J Anaesthesiol 2003 ; 20: A-359.
23. Effect of remifentanil on Sevoflurane MACBAR in women.  
Anesthesiology 2004; 101: A-474.
24. Effect site concentration of remifentanil (CE50) blunting cardiovascular responses to tracheal intubation and skin incision during BIS-guided anesthesia.  
Anesthesiology 2004; 101: A-476.
25. The effect of adding nitrous oxide on MACBAR of sevoflurane combined with two target-controlled (1-3 ng/ml) of remifentanil in women.  
Anesthesiology 2004; 101: A-475
26. Effect of two target-controlled concentrations (1-3 ng/ml) of remifentanil on MACBAR of desflurane.  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005.
27. A new approach to the PK-PD of remifentanil  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
28. Cerebral state index vs bispectral index as electroencephalogram measures of anesthetic drug effect during propofol anesthesia.  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
29. Predictive performance of “Diprifusor” TCI system in morbidly obese patients during open bariatric surgery  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
30. Sedation for upper gastrointestinal tract ultrasound endoscopy: a comparison of anaesthetist administered midazolam-phetidine and patient controlled sedation with propofol-fentanyl.  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
31. The effect of metachromatic leukodystrophy (MLD) on TPS and propofol requirements in children undergoing magnetic resonance imaging (MRI)  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
32. Acute Pain Service and multimodal therapy for postsurgical pain control: evaluation of side effects.  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
33. Acute Pain Service and multimodal therapy for postsurgical pain control: cost evaluation.  
SIAARTI Bari 2005 Minerva Anesthesiol vol 71 suppl 2 n 10 Ottobre 2005
34. A new approach to the PK-PD of remifentanil.  
Anesthesiology 2005; 103: A-775.
35. Predictive performance of “Diprifusor” TCI system in morbidly obese patients during open bariatric surgery under propofol/remifentanil anesthesia.

Anesthesiology 2005; 103: A-773.

36. Effect site concentrations of remifentanil blunting cardiovascular responses to surgical stimuli during BIS guided propofol anesthesia in morbidly obese patients. Anesthesiology 2005; 103: A-774.

37. Effect of two target-controlled concentrations (1-3 ng/ml) of remifentanil on MACBAR of desflurane. Anesthesiology 2005; 103: A-772.

38. Sedation for upper gastrointestinal tract ultrasound endoscopy: a comparison of anaesthetist administered midazolam-phetidine and patient controlled sedation with propofol-fentanyl. Anesthesiology 2005; 103: A-621

39. Perioperative management of a patient with Takayasu's arteritis admitted for caesarean section. Anesthesiology 2006; 105: A-1670.

40. Effect of obesity on wash-in and wash-out curves of desflurane. Anesthesiology 2006; 105: A-636.

41. Wash-in and wash-out curves of sevoflurane and desflurane in morbidly obese patients. Anesthesiology 2006; 105: A-639.

42. Comparison of CSM and BIS monitoring during EUS of upper gastrointestinal tract. Eur J Anaesthesiol 2006 ; 23: A-106

43. MACBAR of desflurane with two target controlled concentrations of remifentanil Eur J Anaesthesiol 2006 ; 23: A-520.

44. A new approach to remifentanil PK-PD Eur J Anaesthesiol 2006 ; 23: A-521

45. Predictive performance of "Diprifusor" TCI in obese patients Eur J Anaesthesiol 2006 ; 23: A-522

46. Total intravenous anesthesia in morbidly obese patients Eur J Anaesthesiol 2006 ; 23: A-561

47. Sedation in paediatric patients affected by metachromatic leukodystrophy. Eur J Anaesthesiol 2006 ; 23: A-626

48. Prospective randomized comparison between kinetic profiles of desflurane and sevoflurane in morbidly obese patients. SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006

49. The effect of adding nitrous oxide on MAC of desflurane combined with two target-controlled concentrations of remifentanil. SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006

50. Analysis of the pharmacokinetic-pharmacodynamic properties of sevoflurane using two depth of anesthesia monitors: the bispectral index and the cerebral state monitor. SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006.

51. CSM vs BIS in corso di anestesia pediatrica con sevoflurane. SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006

52. CSM vs BIS in corso di anestesia pediatrica con propofol. SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006

53. Application of bispectral index and cerebral state index to the measurement of the

- electroencephalographic effect of sevoflurane.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
54. Può l'analisi di alcune covariate avere un valore predittivo sull'esito della NIMV condotta in ambiente non intensivo?  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
55. La NIMV nell'insufficienza respiratoria cronica riacutizzata: analisi retrospettiva di 2 anni di attività in reparto non-intensivo.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
56. La NIMV nella polmonite: analisi retrospettiva di 2 anni di attività in reparto non-intensivo.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
57. The bispectral index monitor: a new tool for aircare helicopter medical personnel.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
58. The effect of morbid obesità on kinetic of desflurane: wash-in wash out curves and recovery times.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
59. Prospective randomized comparison between spinal and general anesthesia on haematological factors affectng outcome during radical retropubic prostatectomy (RRP).  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
60. La NIMV nell'edema polmonare acuto: analisi retrospettiva di 2 anni di attività in reparto non-intensivo.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
61. Continuous epidural analgesia in laparotomic major upper surgery for postoperative pain control.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
62. Continuous epidural analgesia and patient controlled analgesia in laparoscopic colorectal surgery for postoperative pain control.  
SIAARTI Bastia Umbra 2006 Minerva Anesthesiol vol 72 suppl 2 n 10 Ottobre 2006
63. Prospective randomized comparison between spinal and general anesthesia on haematological factors affectng outcome during radical retropubic prostatectomy (RRP).  
Eur J Anaesthesiol 2007 ; 24: 6AP3-2
64. Effect of nitrous oxide on the desflurane requirement for blunting sympathetic responses after surgical incision combined with two target-controlled concentrations of remifentanil (1 and 3 ng/ml)  
Eur J Anaesthesiol 2007 ; 24: 9AP3-2
65. How to correct Minto's pharmacokinetic set in morbidly obese patients: the "fictitious height" in women (II).  
1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.
66. How to correct Minto's pharmacokinetic set in morbidly obese patients: the "fictitious height" in men (II).  
1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.
67. How to correct Minto's pharmacokinetic set in morbidly obese patients: the "fictitious height" in women (I).  
1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.
68. How to correct Minto's pharmacokinetic set in morbidly obese patients: the "fictitious height" in men (I).

1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.

69. Greater peripheral blood flow but less bleeding with propofol vs sevoflurane during spine surgery.

1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.

70. Weight input for Target Controlled Infusion of remifentanil in morbidly obese patients: testing a possible correction.

1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.

71. Target Controlled Infusion of remifentanil in morbidly obese patients: pursuing the best weight input correction.

1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy

## Reports to Congress = 17

1. Anestesia nel grande anziano  
SIAARTI Napoli 2000
2. New drugs for the anesthesiologist: practical effects on daily practice. Atti del Convegno Clinical Practice and Pain Management in Orthopedic Anesthesia. Milano, 2-4 March, 2000.  
  
Unilateral spinal block: technical issues. Highlights in Regional Anaesthesia and Pain Therapy IX. 2000:208-11.
3. ED50 of ropivacaine and bupivacaine in a model of peripheral nerve block. International Monitor Regional Anesthesia Pain Medicine 2000;12:A17.
4. La PCA epidurale  
SIAARTI Milano 2002
5. Quale tecnica in day surgery: ottimizzazione delle tecniche  
SIAARTI Milano 2002
6. TCI e BIS  
SIAARTI Milano 2002
7. Come utilizzare gli ipnotici  
Foundation Congress of Italian SIVA. Roma 2002
8. Come utilizzo gli analgesici nella TIVA  
SIAARTI Roma 2003
9. Sistemi di somministrazione di Remifentanil: nuovi orizzonti  
SIAARTI Roma 2003
10. Monitored Anesthesia Care  
SIAARTI Roma 2003
11. Recupero Intraoperatorio della coscienza  
SIAARTI Genova 2004
12. Gli oppioidi a confronto con gli anestetici alogenati di ultima generazione.  
SIAARTI Bari 2005
13. Farmacocinetica-farmacodinamica nei pazienti grandi obesi.  
SIAARTI Bari 2005
14. Esperienza gestionale di un servizio di Acute Pain Service  
SIAARTI Bari 2005
15. Anestesia generale bilanciata nel paziente obeso.  
SIAARTI Bastia Umbra 2006
16. TIVA-TCI in morbidly obese patients.  
EuroSiva Advanced Course in TIVA-TCI 2006, Ancona, Italy
17. Servin formula vs no correction for weight input in TCI devices during propofol-remifentanil TCI during bariatric surgery.  
1<sup>st</sup> World Congress of Total Intravenous Anaesthesia – TCI, 2007, Venice Italy.



## Research Grants

1. Sevoflurane pharmacokinetics in morbidly obese patients. **1st National prize** for the best communication on Anaesthesia, Pain Management and Hyperbaric Therapy.  
54° National Congress SIAARTI.
2. Ropivacaine and bupivacaine equipotency in peripheral nerve blocks. **2nd National prize** for the best communication related to topics of regional anaesthesia  
54° National Congress SIAARTI.
3. The effect of remifentanil on sevoflurane requirements for somatic and sympathetic responses to surgical incision. **1st National prize** for the best communication on Anaesthesia, Pain Management and Hyperbaric.  
56° National Congress SIAARTI
4. The effect of remifentanil on sevoflurane requirements for sympathetic responses to surgical incision in women. **1st National prize** for the best communication on Anaesthesia, Pain Management and Hyperbaric..  
57° National Congress SIAARTI.

## Articles on medical magazines = 4

1. "Un bisturi che lascia il segno" Medici Oggi n.2 Marzo 2000
2. "Vincere la sofferenza" Medici Oggi n.6 Settembre 2000
3. "Anestesia bilanciata versus Total Intravenous Anesthesia (TIVA): uso, vantaggi e prospettive" Perimed – medicina perioperatoria 2008; vol 2 – n°1.
4. "Farmaci equivalenti in sala operatoria" Perimed – medicina perioperatoria 2010; Suppl n° 1


## Books = 6

1. Coauthor of "Dolore postoperatorio: la teoria e la pratica" –M. Berti. CE.DI.S. Edition, 1999.
2. Author of "Remifentanil – applicazioni pratiche e associazioni terapeutiche" Ed. Interaction, Milan, 2000.
3. Author of "Monitored Anesthesia Care – dalla sedazione alla MAC. Una guida pratica" Ed. UTET Milan 2002
4. Coauthor of "Awareness in anestesia" –G. Torri. Minerva Medica Edizioni, 2004
5. Coauthor of "Il dolore acuto postoperatorio" –G. Torri. Minerva Medica Edizioni, 2006
6. Coauthor of "Chirurgia e riabilitazione della mano".-G. Pajardi. PICCIN 2012

Il sottoscritto è a conoscenza che, ai sensi dell'art. art. 76 del DPR 445/2000, le dichiarazioni mendaci, la falsità negli atti e l'uso di atti falsi sono puniti ai sensi del codice penale e delle leggi speciali. Inoltre, il sottoscritto autorizza al trattamento dei dati personali ai sensi dell'art. 13 D. Lgs. 30 giugno 2003 n°196 – "Codice in materia di protezione dei dati personali" e dell'art. 13 GDPR 679/16 – "Regolamento europeo sulla protezione dei dati personali".

22/03/2021

Albertin Andrea

Firma..... 

M.D. Ph.

